

MYAH

Optical Biometry + Corneal Topography

Build, Manage and Grow
Your Myopia & Dry Eye Practice



LEARN MORE at
topconmyah.com

 **TOPCON** Healthcare

Myopia greatly impacts the quality of life and personal development of children¹.

It has never been a better time to join the battle against the global myopia epidemic. MYAH is the perfect instrument for eyecare professionals interested in building, managing and growing a myopia service.

Overview of MYAH



Corneal Topography including keratoconus screening and pupillometry



Axial length measurement using optical low coherence Interferometry



Progression reports for analysing treatment efficacy



Myopia and dry eye questionnaires



Comprehensive suite of dry eye assessment tools



Patient-friendly with rapid capture



Compact, space-saving, easy to operate



Exclusive axial length reference databases

Did you know that 50% of the world’s population² may be myopic by 2050?

European regions are not an exception.

Region	2000	2010	2020	2030	2040	2050
Central Europe	20.5%	27.1%	34.6%	41.8%	48.9%	54.1%
Eastern Europe	18.0%	25.0%	32.2%	38.9%	45.9%	50.4%
Western Europe	21.9%	28.5%	36.7%	44.5%	51.0%	56.2%
Global	22.9%	28.3%	33.9%	39.9%	45.2%	49.8%

Building a myopia management practice requires you to educate your patients and their families about the implications of myopia progression, to manage the condition and to grow your service offering.

BUILD YOUR MYOPIA MANAGEMENT SERVICE

MYAH establishes the initial baseline to monitor risk, allowing you to start the conversation early with parents.

MANAGE: MONITOR AND COMPARE

MYAH provides essential information to monitor eye elongation and compare axial length measurements with built-in growth curves.

GROW YOUR MYOPIA MANAGEMENT SERVICE

Offering axial length screening tests may complement your refraction tests.

MYAH offers all the technologies required to support myopia management: optical biometry, corneal topography and pupillometry — it is a one-time investment. In addition, MYAH is an all-in-one device that offers an evolving platform which provides the tools to add or grow a dry-eye management service.




Introducing MYAH's growth curves.

MYAH allows you to monitor the progression of myopia and compare measurements with the growth curves for axial length.


The majority of myopic eyes become myopic principally because of excessive axial elongation³. By using the extensive axial length dataset collected by Erasmus University (Rotterdam, NL)⁴, or the dataset on Chinese children (Shanghai)^{5,6}, now incorporated into MYAH, you can monitor axial length and then compare the patient's data with normative growth curves. You will thus be able to better understand a child's risk of myopia in adulthood.

You can now enhance your myopia management service with MYAH's growth curves.



New! Questionnaires on myopia & dry eye


Discover the latest additions to MYAH's services: a new myopia questionnaire along with two comprehensive dry eye questionnaires: OSDI and DEQ-5. These insightful questionnaires help systematically track data over time and are seamlessly integrated into our enhanced reports.



Additional Features.

Dynamic Pupillometry


Provides clear information on the reaction time and size of the pupil, which may be useful to monitor low dose atropine compliance or to titrate the dose of atropine. The user can examine pupil centration and diameter over a range of light levels, which is useful for Ortho-K and multifocal lens fittings, and is also informative for pre- and post-refractive surgery.



Contact Lens Fitting

MYAH provides support for contact lens fitting, reducing the number of lenses that need to be trialed on the eye:

- Includes a database of conventional RGP and Ortho-K lenses.
- Export topography data to 3rd party calculators.
- Fluorescein simulation with ability to save and review data.



Dry Eye Assessment Tools

These tools offer non-invasive tear break-up time (NIBUT), Meibomian gland imaging with the area of loss analysis, tear meniscus height analysis, blink analysis, real fluorescein imaging and video acquisition. You can play a video of the topographic maps and also the Zernike maps and see them change during the break up of the tear film. Review results, print and export reports on network or USB.



Corneal Topography


MYAH offers another range of tools to analyse the anterior cornea, including topographic maps, 3D maps, comparison maps, height maps, Zernike analysis and keratoconus screening.



Corneal Aberration Summary

The Zernike expansion coefficient is used to determine which component(s) dominate the aberration structure of the cornea and to what degree.

The anterior corneal Zernike summary consists of 36 polynomials up to the 7th order and provides a clear view of the optical irregularities that can impact the quality of vision.



Topcon MYAH

The all-in-one instrument features tools that monitor eye elongation, assess dry eye and assist with contact lens fittings.



MYAH makes your practice dynamic and smart.

This versatile instrument, with its intuitive and user-friendly interface, integrates easily into your workflow and offers different options for exporting the results.

4 EASY STEPS



Select patient** and acquisition mode.



Align patient and adjust automated chinrest.



Follow alignment guides to focus and trigger to start.



Review results and print/export reports to network or USB.

** Create new patient, select existing patient or select patient from DICOM (search/worklist).

MYAH SPECIFICATION

FEATURE	SPECIFICATION
Keratoscopic cone	24 rings equally distributed on a 43 D sphere
Analyzed points	Over 100,000
Measured points	6,144
Corneal coverage	Up to 9.8 mm on a sphere of radius 8.00 mm (42.2 D with n=1.3375)
Axial Biometry	Low-coherence interferometry on optical fiber (SLED @ 820 nm)
Capture system	Guided-focus
Database	Internal
Pupillometry	Dynamic, Photopic, Mesopic, Scotopic
Fluorescein	Image, Video
Reports	Keratometry, comparison map, contact lens, height map, Zernike analysis, pupillometry, Meibomian glands, tear film break-up time, tear meniscus height, Rx/AL trend analysis, fluorescein, parental, dry eye summary & dry eye follow-up report
Working environment	10 °C – 40 °C, Relative humidity 8 – 75% (no condensing), Atmospheric pressure 800 – 1060 hPa
Power supply	AC 100 – 240 V 50/60 Hz
Power consumption	100 VA
Dimensions	320 mm (W) x 490 mm (H) x 470 mm (L), 18 Kg
Printing options	USB printer, Network printer, PDF on network shared folder, PDF on USB PDF or Image on network folder or on USB
Operating System	Windows 10 64-bit
Monitor	LCD 10.1 inch capacitive touch screen
RAM	At least 4 GB
Hard Disk	500 GB
External connections	LAN integrated, 2x USB

INFORMATION ON MEASUREMENTS

MEASUREMENT	MEASURING RANGE	DISPLAY RESOLUTION	IN VIVO REPEATABILITY
Keratometry	Radius of curvature	5.00 – 12.00 mm	±0.02 mm
	Curve Radius in Diopter (D) (n=1,3375)	28.00 – 67.50 D	±0.12 D
Axial Length	15.00 – 36.00 mm	0.01 mm	±0.027 mm
Pupil dimension	0.50 – 10.00 mm	0.01 mm	N/A
Limbus (White-To-White)	8.00 – 14.00 mm	0.01 mm	±0.05 mm
IBI Index (Interblink Interval)	0.2 – 20.0 s	0.1 s	N/A
Non-invasive Break-Up Time (TBT)	0.5 – 30.0 s	0.1 s	N/A
Meibomian Glands area of loss	0 – 100%	1%	N/A
Tear Meniscus Height	0.10 – 1.00 mm	0.01 mm	N/A

Not all products, services or offers are approved or offered in every market, and products vary from one country to another. Contact your local distributor for country-specific information.

1. Report of the Joint World Health Organization-Brien Holden Vision Institute. Global Scientific Meeting on Myopia. The Impact of myopia and high myopia. University of New South Wales, Sydney, Australia. 16-18 March 2015.

2. Holden, BA, Fricke, TR, Wilson, DA et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. Ophthalmology. 2016; 123:1036-42. Available from: doi: 10.1016/j.ophtha.2016.01.006

3. (Gifford KL, Richdale K, Kang P, Aller TA, Lam CS, Liu YM, Michaud L, Mulder J, Orr JB, Rose KA, Saunders KJ, Seidel D, Tideman JWL, Sankaridurg P. IMI - Clinical Management Guidelines Report. Invest Ophthalmol Vis Sci. 2019 Feb 28;60(3):M184-M203).

4. Coordinates incorporated in this Myopia device are the most recent available data and originate from the Myopia Research Group of Erasmus MC, Rotterdam

5. Courtesy of Prof. Xu Xun, MD

6. He X, Sankaridurg P, Naduvilath T, Wang J, Xiong S, Weng R, Du L, Chen J, Zou H, Xu X. Normative data and percentile curves for axial length and axial length/corneal curvature in Chinese children and adolescents aged 4-18 years. Br J Ophthalmol. 2023 Feb;107(2):167-175

IMPORTANT

Subject to change in design and/or specifications without advanced notice. In order to obtain the best results with this instrument, please be sure to review all user instructions prior to operation. Medical device MDR Class IIa. Manufacturer: VISIA imaging S.r.l.

TOPCON CORPORATION

75-1 Hasunuma-cho, Itabashi-ku, Tokyo 174-8580, JAPAN.
Phone: +81-(0)3-3558-2522/2502 Fax: +81-(0)3-3965-6898
www.topconhealthcare.jp

VISIA IMAGING S.R.L.

Via Martiri della Libertà 95/e, 52027 San Giovanni Valdarno (AR), ITALY



TOPCON HEALTHCARE SOLUTIONS EMEA OY

Saaristonkatu 9, 90100 Oulu, FINLAND
Phone: +358-20-734-8190
E-mail: thsemea.sales@topcon.com
www.topconhealthcare.eu

TOPCON SCANDINAVIA

Neogatan 2, P.O.Box 25, 43151 Mölndal
SWEDEN
Phone: +46-(0)31-7109200
Fax: +46-(0)31-7109249
E-mail: info.hcs.se@topcon.com
www.topconhealthcare.eu/sv_SE

TOPCON FRANCE MEDICAL

1 rue des Vergers, Parc Swen,
Bâtiment 2, 69760 Limonest, FRANCE
Phone: +33-(0)4-37 58 19 40
Fax: +33-(0)4-72 23 86 60
E-mail: topconfrance@topcon.com
www.topconhealthcare.eu/fr_FR

TOPCON GREAT BRITAIN MEDICAL

Topcon House, Kennet Side, Bone Lane,
Newbury, Berkshire RG14 5PX
UNITED KINGDOM
Phone: +44-(0)1635-551120
Fax: +44-(0)1635-551170
E-mail: medical.tgbm@topcon.com
www.topconhealthcare.eu/en_UK

TOPCON EUROPE MEDICAL B.V.

Essebaan 11, 2908 LJ Capelle a/d IJssel
P.O. Box 145, 2900 AC Capelle a/d IJssel
THE NETHERLANDS
Phone: +31-(0)10-4585077
Fax: +31-(0)10-4585045
E-mail: medical@topcon.com
www.topconhealthcare.eu

TOPCON ESPAÑA

Oficina Principal en España
Frederic Mompou, 4,
08960 Sant Just Desvern, Barcelona, SPAIN
Phone: +34-93-4734057
Fax: +34-93-4733932
E-mail: medica@topcon.com
www.topconhealthcare.eu/es_ES

TOPCON DEUTSCHLAND MEDICAL

Hanns-Martin-Schleyer Strasse 41,
D-47877 Willich, GERMANY
Phone: (+49)2154-885-0
Fax: (+49)2154-885-177
E-mail: info@topcon-medical.de
www.topconhealthcare.eu/de_DE

TOPCON IRELAND MEDICAL

Unit 292, Block G, Blanchardstown,
Corporate Park 2 Ballycoolin
Dublin 15, D15 DX58, IRELAND
Phone: +353-12233280
E-mail: medical.ie@topcon.com
www.topconhealthcare.eu/en_IE

TOPCON DANMARK

Praestemarksvej 25, 4000 Roskilde
DANMARK
Phone: +45-46-327500
Fax: +45-46-327555
E-mail: info.todk@topcon.com
www.topconhealthcare.eu/dk_DK

TOPCON ITALY

Viale dell' Industria 60,
20037 Paderno Dugnano, (MI) ITALY
Phone: +39-02-9186671
Fax: +39-02-91081091
E-mail: info@topcon.com
www.topconhealthcare.eu/it_IT

TOPCON POLSKA SP. Z O.O.

ul. Warszawska 23, 42-470 Siewierz
POLAND
Phone: +48-(0)32-670-50-45
Fax: +48-(0)32-671-34-05
E-mail: info.tpl@topcon.com
www.topconhealthcare.eu/pl_PL

medical@topcon.com | topconmyah.com
M000092E-3

TOPCON Healthcare